arXiv:1401.6429v1 [math.DG] 10 Dec 2013

ALMOST CONTACT CURVES IN TRANS-SASAKIAN 3-MANIFOLDS

S. K. Srivastava[1]
Department of Mathematics, Central University of Himachal Pradesh,
Dharamshala - 176215, Himachal Pradesh, India

Abstract

This paper is devoted to the study of curvature and torsion of almost contact curves in trans-Sasakian 3-Manifolds. The conditions for the frenet curves to be almost contact curves in trans-Sasakian 3-manifolds have been obtained.

AMS Subject Classification 2010: 53D15, 53C25.
Key words: Almost contact curve, Frenet curve, trans- Sasakian manifold.

1 Introduction

Almost contact curves play a important role in geometry and topology of contact metric manifolds, e.g. *a diffeomorphism of a contact manifold is a contact transformation if and only if any almost contact curves in a domain of it go to almost contact curves* [5]. Several authors have studied almost contact curves in contact geometry such as [1, 5, 6, 9, 11, 12, 16, 18]. In [5] Baikoussis and Blair have studied almost contact curves in contact metric 3-manifold and gave the Frenet 3-frame in this class of manifold. Belkhelfa *et al.* have extended some of the results of [5] from the Riemannian to the Lorentzian case, and classified all biharmonic almost contac curves in Sasaki-Heisenberg spaces[18]. A result of Blair [see theorem 8.2, p.134 [1]] had been generalized by Welyczko[11] to the case of 3-dimensional Quasi-Sasakian manifolds. Moreover the author had also obtained some interesting properties of non-Frenet almost contact curves in normal almost paracontact metric 3-manifolds [12]. Özgür and Tripathi established necessary and sufficient conditions for almost contact curves having parallel mean curvature vector, proper mean curvature vector, being harmonic and being of type $AW(k), k = 1, 2, 3$; in α- Sasakian manifolds[6]. In [9] Lee characterized almost contact curves in a Sasakian manifold having the following properties: (i) a pseudo-Hermitian parallel mean curvature vector field (ii) a pseudo-Hermitian proper mean curvature vector field in the normal bundle. Recently, Inoguchi and Lee have studied almost contact curves in normal almost contact metric 3-manifold satisfying $\nabla H = \lambda H$ or $\nabla^{\perp} H = \lambda H$ and gave natural equations for *planar biminimal* curves [16].

The purpose of this paper is to investigate the properties of almost contact curves in trans-Sasakian 3-manifolds. This paper is organized as follows: In §2 we recall some basic definitions and facts about almost contact metric (in brief a.c.m.) manifolds, trans-Sasakian manifolds and Frenet curves. The curvature, torsion of almost contact curves and the conditions for the frenet curves to be almost contact curves in trans-Sasakian 3-manifolds have been obtained, and finally we construct the examples in §3.

[1]Email: sachink.ddumath@gmail.com

2 Preliminaries

2.1 Contact metric manifolds

A $(2n+1)$-dimensional differentiable manifold M is said to be an *almost contact manifold* if its structural group $GL_{2n+1}\mathbb{R}$ of linear frame bundle is reducible to $\mathrm{U}(n) \times \{1\}$ (Gray [10]). This is equivalent to existence of a tensor field of type (1, 1), a vector field ξ and a 1-form η satisfying

$$\phi^2 = -I + \eta \otimes \xi, \qquad \eta(\xi) = 1 \tag{2.1}$$

From these conditions one can easily obtain

$$\phi\xi = 0, \qquad \eta o \phi = 0. \tag{2.2}$$

Moreover, scince $\mathrm{U}(n) \times \{1\} \subset \mathrm{SO}(2n+1)$, M admits a Riemannian metric g satisfying

$$g(\phi X, \phi Y) \;=\; g(X,Y) - \eta(X)\eta(Y) \tag{2.3}$$

for all $X, Y \in \Gamma(TM)$. Such a metric is called an *associated metric* (Sasaki [19]) of the almost contact manifold M. With respect to g, η is metrically dual to ξ, that is

$$g(X,\xi) \;=\; \eta(X) \tag{2.4}$$

A structure (ϕ, ξ, η, g) on M is called an *almost contact metric structure* and a manifold M equipped with an almost contact metric structure is said to be an *almost contact metric manifold*. The fundamental 2-form Φ of the manifold is defined by

$$\Phi(X,Y) = g(X, \phi Y) \tag{2.5}$$

for all $X, Y \in \Gamma(TM)$. An almost contact metric manifold M is said to be a *contact metric manifold* if $\Phi = d\eta$. Here the exterior derivative $d\eta$ is defined by

$$d\eta(X,Y) = \frac{1}{2}\left(X\eta(Y) - Y\eta(X) - \eta([X,Y])\right). \tag{2.6}$$

On a contact metric manifold, η is *contact form*, i.e., $\eta \wedge (d\eta)^n \neq 0$ everywhere on M. In particular, $\eta \wedge (d\eta)^n \neq 0$ is a volume element on M so that a contact manifold is orientable. Define a (1, 1) type tensor field h and l by $h = \frac{1}{2}L_\xi \phi,\ lX = R(., \xi)\xi$, where L denotes the Lie differentiation and R the curvature tensor respectively. The operators h and l are self-adjoint and satisfy: $h\xi = l\xi = 0$ and $h\phi = -\phi h$. Also we have $Tr.h = Tr.\phi h = 0$. Moreover, if ∇ denotes the Levi-Civita connection on M, then following formulas holds on a contact metric manifold.

$$\nabla_X \xi = -\phi X - \phi h X. \tag{2.7}$$

$$l = \phi l \phi - 2(h^2 + \phi^2). \tag{2.8}$$

On the direct product manifold $M \times \mathbb{R}$ of an almost contact metric manifold M and the real line $\mathbb{R}$, any tangent vector field can be represented as the form $\left(X, f\frac{d}{dt}\right)$, where $X \in \Gamma(TM)$ and f is a function on $M \times \mathbb{R}$ and t is the cartesian coordinate on the real line $\mathbb{R}$.
Define an almost complex structure J on $M \times \mathbb{R}$ by

$$J\left(X, \lambda\frac{d}{dt}\right) = \left(\phi X - \lambda\xi, \eta(X)\frac{d}{dt}\right). \tag{2.9}$$

If J is integrable then M is said to be *normal*. Equivalently, M is normal if and only if

$$[\phi, \phi](X, Y) + 2d\eta(X, Y)\xi = 0, \tag{2.10}$$

where $[\phi, \phi]$ is the Nijenhuis torsion tensor of ϕ defined by

$$[\phi, \phi](X, Y) = [\phi X, \phi Y] + \phi^2[X, Y] - \phi[\phi X, Y] - \phi[X, \phi Y] \tag{2.11}$$

for all $X, Y \in \Gamma(TM)$.
For an arbitrary almost contact metric 3-manifold M, we have ([20]):

$$(\nabla_X\phi)Y = g(\phi\nabla_X\xi, Y)\xi - \eta(Y)\phi\nabla_X\xi, X \in \Gamma(TM) \tag{2.12}$$

where ∇ is the Levi-Civita connection on M.

2.2 trans-Sasakian manifolds

This class of manifolds arose in a natural way from the classification of almost contact metric structures and they appear as a natural generalization of both Sasakian and Kenmotsu manifolds. In [4] Gray Harvella classification of almost Hermite manifolds appear as a class W_4 of Hermitian manifolds which are closely related to locally conformally Kähler manifolds. An almost contact metric structure on a manifold $M \times \mathbb{R}$ belongs to the class W_4. The class $C_6 \oplus C_5$ [13] coincides with the class of trans-Sasakian structure of type (α, β).

An almost contact metric structure (ϕ, ξ, η) on a connected manifold M is called *trans- Sasakian structure* [14] if $(M \times \mathbb{R}, J, G)$ belongs to the class W_4 [4], where J is the almost complex structure defined by (2.9) and G is the product metric on $M \times \mathbb{R}$. This may be expresses by the condition [3]

$$(\nabla_X\phi)Y = \alpha(g(X, Y)\xi - \eta(Y)X) + \beta(g(\phi X, Y)\xi - \eta(Y)\phi X) \tag{2.13}$$

for the smooth functions α and β on M. Hence we say that the trans - Sasakian structure is of type (α, β). From (2.13) it follows that

$$(\nabla_X\eta)(Y) = -\alpha g(\phi X, Y) + \beta g(\phi X, \phi Y) \tag{2.14}$$

We note that trans-Sasakian structure of type $(0, 0)$, $(\alpha, 0)$ and $(0, \beta)$ are the *cosymplectic*, $\alpha-$Sasakian and $\beta-$Kenmotsu manifold respectively.

2.3 Frenet Curves

Let (M, g) be a Riemannian $n-$manifold with Levi-Civita connection ∇. A unit speed curve $\gamma : I \to M$ is said to be an $r-$*Frenet curve* [1] if there exists an orthonormal $r-$frame field $(E_1 = \gamma', E_2, ..., E_r)$ along γ such that there exist positive smooth functions $k_1, k_2, ..., k_{r-1}$ satisfying

$$\nabla_{\gamma'} E_1 = k_1 E_2, \quad \nabla_{\gamma'} E_2 = -k_1 E_1 + k_2 E_3, ..., \nabla_{\gamma'} E_r = -k_{r-1} E_{r-1}.$$

The function k_r is called the $r-$th curvature of γ. A Frenet curve is said to be

• a *geodesic* if $r = 1$, i.e., $\nabla_{\gamma'} \gamma' = 0$.

• a *Riemannian circle* if $r = 2$, and k_1 is non-zero constant.

• a *helix* of order r if $k_1, k_2, ..., k_{r-1}$ are constants.

In case $n = 3$, we denote by $(E_1, E_2, E_3) = (T, N, B)$. Then we have the Serret-Frenet equation:

$$\nabla_T T = kN, \quad \nabla_T N = -kT + \tau B \text{ and } \nabla_T B = -\tau N \tag{2.15}$$

where $T = \gamma'$.

The first curvature $k = k_1$ and the second curvature $\tau = k_2$ are called the *geodesic curvature* and *geodesic torsion* of γ, respectively. The vector field N and B are called the unit normal vector field and binormal vector field of γ, respectively.

3 Almost contact Curves

Let $\gamma : I \to M$ be a curve parameterized by arc-length (the *natural parametrization*) in an almost contact metric 3-manifold M with Frenet frame (T, N, B).

Definition 3.1 *A Frenet curve γ in an almost contact 3-manifold M is said to be an almost contact curve if it is an integral curve of the contact distribution $D = ker\eta$, equivalently, $\eta(\gamma') = 0$.*

In particular, when $\eta \wedge (d\eta)^n \neq 0$, almost contact curves are traditionally called *Legendre curves* (cf.[5]). We begin with a proposition that will motivate the main result:

Proposition 3.2 *Let M be a trans-Sasakian 3-manifold. Then for non-geodesic almost contact curve $\gamma : I \to M$, curvature (κ) and torsion (τ) are given by*

$$\kappa = \sqrt{\beta^2 + \vartheta^2} \tag{3.1}$$

$$\tau = |\alpha + \frac{\beta\vartheta' - \beta'\vartheta}{\kappa^2}| \tag{3.2}$$

Proof: Let γ be an almost contact curve on M. Then

$$\nabla_{\gamma'} T = \nabla_{\gamma'} \gamma' = -\beta\xi + \vartheta\phi\gamma' \tag{3.3}$$

for some function ϑ. The unit normal vector field N is given by

$$N = \frac{1}{\kappa} \nabla_{\gamma'} T = -\frac{\beta}{\kappa}\xi + \frac{\vartheta}{\kappa}\phi\gamma' \tag{3.4}$$

Differentiating (3.4) along γ', we get

$$\nabla_{\gamma'} N = -\kappa\gamma' + p\xi + q\phi\gamma' \tag{3.5}$$

where

$$p = \frac{\vartheta}{\kappa}\alpha - \frac{\beta'\kappa - \beta\kappa'}{\kappa^2}, \qquad q = \frac{\alpha\beta}{\kappa} + \frac{\vartheta'\kappa - \vartheta\kappa'}{\kappa^2}.$$

Here β', δ' and κ' are

$$\beta'(s) = \frac{d}{ds}\beta(\gamma(s)), \vartheta'(s) = \frac{d}{ds}\vartheta(\gamma(s)) \text{ and } \kappa'(s) = \frac{d}{ds}\kappa(\gamma(s)).$$

From (3.4) and $\tau B = \nabla_{\gamma'} N + \kappa T = p\xi + q\phi\gamma'$, we have (3.1) and (3.2). □

MAIN RESULT

Theorem 3.3 *For a Frenet curve $\gamma : I \to M$ in a trans-Sasakian 3-manifold M with $\alpha \neq 0$ and $\beta \neq 0$. Set $\sigma = \eta(\gamma')$. If $\tau = |l_1\alpha + l_2\beta + l_3|$ and at one point of I, $\sigma = \sigma' = \sigma'' = 0$, then γ is an almost contact curve.*

Where

$$l_1 = \frac{1}{\sqrt{1-\sigma^2}}, \ \ l_2 = -\frac{pq\sigma}{\sqrt{1-\sigma^2}(p^2+q^2)}, \ \ l_3 = -\frac{p^2}{(p^2+q^2)}\gamma\left(\frac{\beta}{p}\right),$$

p is non - zero constant on I and q is certain function on I.

Proof: Suppose that γ' is not collinear with ξ and describe curvature (κ) and torsion (τ) of γ on I. We may decompose $\nabla_{\gamma'}\gamma'$ as

$$\nabla_{\gamma'}\gamma' = \nabla_{\gamma'} T = \frac{p}{\sqrt{1-\sigma^2}}\phi\gamma' + \frac{q}{\sqrt{1-\sigma^2}}\left(\xi - \sigma\gamma'\right) \tag{3.6}$$

Therefore

$$k = \sqrt{p^2+q^2} \tag{3.7}$$

is curvature of γ.

Using (2.12) and (3.6), we have

$$\begin{aligned} \sigma' &= \gamma'\left(g\left(\xi,\gamma'\right)\right) \\ &= g\left(\nabla_{\gamma'}\xi,\gamma'\right) + g\left(\xi,\nabla_{\gamma'}\gamma'\right) \\ &= \beta\left(1-\sigma^2\right) + q\sqrt{1-\sigma^2}. \end{aligned} \tag{3.8}$$

From (3.6), we find

$$\begin{aligned} N &= \frac{1}{k}\nabla_{\gamma'} T \\ &= \frac{p}{k\sqrt{1-\sigma^2}}\phi\gamma' + \frac{q}{k\sqrt{1-\sigma^2}}\left(\xi - \sigma\gamma'\right). \end{aligned} \tag{3.9}$$

Let us write,

$$p_1 = \frac{p}{k\sqrt{1-\sigma^2}}, q_1 = \frac{q}{k\sqrt{1-\sigma^2}}. \tag{3.10}$$

Then (3.9) becomes

$$N \;\; = \;\; p_1\phi\gamma' + q_1\left(\xi - \sigma\gamma'\right). \tag{3.11}$$

From (3.7) and (3.8), we compute

$$\begin{aligned} p_1' &= \frac{q\left(p'q - pq'\right)}{k^3\sqrt{1-\sigma^2}} + \frac{pq\sigma}{k\left(1-\sigma^2\right)} + \frac{p\sigma\beta}{k\sqrt{1-\sigma^2}}, \\ q_1' &= \frac{p\left(pq' - p'q\right)}{k^3\sqrt{1-\sigma^2}} + \frac{q^2\sigma}{k\left(1-\sigma^2\right)} + \frac{q\sigma\beta}{k\sqrt{1-\sigma^2}}. \end{aligned} \tag{3.12}$$

Differentiating (3.11) along γ', we have

$$\nabla_{\gamma'}N = p_1'\phi\gamma' + p_1\left(\left(\nabla_{\gamma'}\right)\gamma' + \phi\nabla_{\gamma'}\gamma'\right) + q_1'\left(\xi - \sigma\gamma'\right) + q_1\left(\nabla_{\gamma'}\xi - \sigma'\gamma' - \sigma\nabla_{\gamma'}\gamma'\right). \tag{3.13}$$

Using (2.13), (2.14), (3.6), (3.8) and (3.5); we get

$$\begin{aligned} \nabla_{\gamma'}N &= \left[\frac{q\left(p'q - pq'\right)}{k^3\sqrt{1-\sigma^2}} + \frac{pq\sigma}{k\left(1-\sigma^2\right)} + \frac{p\sigma\beta}{k\sqrt{1-\sigma^2}}q\right]\phi\gamma' \\ &+p_1\left[\alpha\left(g\left(\gamma',\gamma'\right)\xi - \eta\left(\gamma'\right)\gamma'\right) + \beta\left(g\left(\phi\gamma',\gamma'\right)\xi - \eta\left(\gamma'\right)\phi\gamma'\right) + \frac{p}{\sqrt{1-\sigma^2}}\left(-\gamma' + \eta\left(\gamma'\right)\xi\right) - \frac{p\sigma}{\sqrt{1-\sigma^2}}\phi\gamma'\right] \\ &+\left[\frac{p\left(pq' - p'q\right)}{k^3\sqrt{1-\sigma^2}} + \frac{q^2\sigma}{k\left(1-\sigma^2\right)} + \frac{q\sigma\beta}{k\sqrt{1-\sigma^2}}\right]\left(\xi - \sigma\gamma'\right) \\ &+q_1\left[-\alpha\phi\gamma' + \beta\left(\gamma' - \sigma\xi\right) - \beta\left(1-\sigma^2\right)\gamma' - q\sqrt{1-\sigma^2}\gamma' - \frac{p\sigma}{\sqrt{1-\sigma^2}}\phi\gamma' - \frac{q\sigma}{\sqrt{1-\sigma^2}}\left(\xi - \sigma\gamma'\right)\right] \end{aligned}$$

which simplifies to,

$$\nabla_{\gamma'}N = \frac{q}{k\sqrt{1-\sigma^2}}\left[\frac{\left(p'q - pq'\right)}{k^2} - \frac{p\sigma}{\sqrt{1-\sigma^2}} - \alpha\right]\phi\gamma' + \frac{p}{k\sqrt{1-\sigma^2}}\left[\frac{\left(pq' - p'q\right)}{k^2} + \frac{p\sigma}{\sqrt{1-\sigma^2}} + \alpha\right]\left(\xi - \sigma\gamma'\right) - k\gamma'$$

or,

$$\nabla_{\gamma'}N = -\frac{q}{k\sqrt{1-\sigma^2}}\left[\frac{\left(pq' - p'q\right)}{k^2} + \frac{p\sigma}{\sqrt{1-\sigma^2}} + \alpha\right]\phi\gamma' + \frac{p}{k\sqrt{1-\sigma^2}}\left[\frac{\left(pq' - p'q\right)}{k^2} + \frac{p\sigma}{\sqrt{1-\sigma^2}} + \alpha\right]\left(\xi - \sigma\gamma'\right) - k\gamma'$$

or,

$$\nabla_{\gamma'}N + k\gamma' = \tau B = p_2\phi\gamma' + q_2\left(\xi - \sigma\gamma'\right) \tag{3.14}$$

where

$$\begin{aligned} p_2 &= -\frac{q}{k\sqrt{1-\sigma^2}}\left[\frac{\left(pq' - p'q\right)}{k^2} + \frac{p\sigma}{\sqrt{1-\sigma^2}} + \alpha\right], \\ q_2 &= \frac{p}{k\sqrt{1-\sigma^2}}\left[\frac{\left(pq' - p'q\right)}{k^2} + \frac{p\sigma}{\sqrt{1-\sigma^2}} + \alpha\right] \end{aligned} \tag{3.15}$$

and

$$p_2^2 + q_2^2 = \frac{1}{(1-\sigma^2)}\left[\frac{(pq' - p'q)}{k^2} + \frac{p\sigma}{\sqrt{1-\sigma^2}} + \alpha\right]^2$$
$$= \tau^2. \tag{3.16}$$

Since, we have assumed that $\tau = |l_1\alpha + l_2\beta + l_3|$. Therefore, we have

$$|l_1\alpha + l_2\beta + l_3| = |\frac{1}{\sqrt{1-\sigma^2}}\left[\frac{(pq' - p'q)}{k^2} + \frac{p\sigma}{\sqrt{1-\sigma^2}} + \alpha\right]|$$

that is,

$$\sigma'' + \frac{2\sigma'^2}{1-\sigma^2} + p^2\sigma = 0. \tag{3.17}$$

For σ not constant, write $\mu = \frac{\sigma'}{p}$. Equation (3.17) yields

$$\mu\frac{d\mu}{d\sigma} + \frac{2\sigma\mu^2}{1-\sigma^2} + \sigma = 0, \tag{3.18}$$

where p is non - zero constant. Integrating (3.18), we have

$$\mu^2 = p^2\left(C\left(1-\sigma^2\right) - 1\right)\left(1-\sigma^2\right) \tag{3.19}$$

where C is contant of integration. Using at one point of I, $\sigma = \sigma' = 0$ and $p \neq 0$ we have $C = 1$. Therefore

$$\sigma'^2 = -p^2\sigma^2(1-\sigma^2).$$

Recalling $\sigma^2 \leq 1$, we have $\sigma = 0$, which is a contradiction. □

Let us suppose that $M = \mathbb{R}^2 \times \mathbb{R}_+$, $\omega : M \to \mathbb{R}_+$ and (x, y, z) be cartesian coordinates in M, we define a trans-Sasakian structure on M by

$$\xi = \frac{\partial}{\partial z}, \;\; \eta = dz - ydx$$

$$\phi = \begin{pmatrix} 0 & -1 & 0 \\ 1 & 0 & 0 \\ 0 & -y & 0 \end{pmatrix},$$

$$g = \begin{pmatrix} \omega + y^2 & 0 & -y \\ 0 & \omega & 0 \\ -y & 0 & 1 \end{pmatrix}.$$

Certain almost contact curves in the above class of manifolds are given below:

Example 3.4. Let us suppose that $\omega = \exp(z)$, then the structure (ϕ, ξ, η, g) is trans-Sasakian structure of type $\left(\frac{-1}{2\exp(z)}, \frac{1}{2}\right)$.

A curve $\gamma = \left(\gamma^1, \gamma^2, \gamma^3\right)$ in M is almost contact curve if and only if

(i) $\dot{\gamma}^3 = \gamma^2 \; \dot{\gamma}^1$

(ii) $\left(\dot{\gamma}^1\right)^2 + \left(\dot{\gamma}^2\right)^2 = \exp\left(-\gamma^3\right)$.

The concrete examples of almost contact curves in M are

(3.4.1) $\gamma(t) = (1, t, 0), t > 0-$ a helix with $\kappa = \tau = 1/2$.

(3.4.2) $\gamma(t) = (lnt, 2, 2lnt), t > 0-$ a curve with $\kappa = 1/2$ and $\tau = 1/2t^2$.

Example 3.5. Let us suppose that $\omega = z$, then the structure (ϕ, ξ, η, g) is trans-Sasakian structure of type $\left(\frac{-1}{2z}, \frac{1}{2z}\right)$.
A curve $\gamma = \left(\gamma^1, \gamma^2, \gamma^3\right)$ *in* M *is almost contact curve if and only if*

(i) $\dot{\gamma}^3 = \gamma^2\ \dot{\gamma}^1$

(ii) $\left(\dot{\gamma}^1\right)^2 + \left(\dot{\gamma}^2\right)^2 = \left(\gamma^3\right)^{-1}$.

The concrete examples of almost contact curves in M are

(3.5.1) $\gamma(t) = (1, t, 0), t > 0-$ a helix with $\kappa = \tau = 1/2$.

(3.5.2) $\gamma(t) = (\sqrt{2t}, \sqrt{2t}, t), t > 0-$ a generalized helix with $\kappa = \tau = 1/2t$.

Acknowledgment: The author wish to express their gratitude to D. E. Blair for helpful comments and remarks in preparing this article.

References

[1] D. E. Blair: *Riemannian Geometry of Contact and Symplectic Manifolds*, Progress in Mathematics 203, Birkhauser Boston, Inc., Boston, MA, 2002.

[2] D. E. Blair, F. Dillen, L. Verstraelen and L. Vrancken: *Deformations of Legendre curves*, Note di Matematica., **15** (1995), no.1, 99-110.

[3] D. E. Blair and J. A. Oubina: *Conformal and related changes of metric on the product of two almost contact metric manifolds*, Publ. Mat., **34** (1990), No. 1, 199-207.

[4] A. Gray and L. M. Hervella: *The sixteen class of almost hermitian manifolds and their linear invariants*, Ann. Mat. Pura Appl. (4) **123** (1980), 35-58.

[5] C. Baikoussis and D. E. Blair: *On Legendre curves in contact 3-manifolds*, Geom. Dedicata, **49** (1994), no. 2, 135-142.

[6] Cihan Özgür and M. M. Tripathi: *On Legendre curves in α- Sasakian manifolds*, Bull. Malays. Math. Sci. Soc. (2) **31** (1)(2008), 91 - 96.

[7] C. Camci, Y. Yayli and H. H. Hacisalihoglu: *On the characterization of spherical curves in 3-dimensional Sasakian spaces*, Jour. Mathematical Analysis and applications, **342** (2008), 1151-1159.

[8] D. Janssens and L. Vanhecke: *Almost contact structures and curvature tensors,* Ködai Math. J. **4** (981), 127.

[9] J. E. Lee: *On Legendre curves in contact Pseudo - Hermitian 3-manifolds,* Bull. Austral. Math. Soc., **81** (2010), 156-164.

[10] J. Gray: *Some global properties of contact structure,* Ann. of Math., **69** (1959), 421-450.

[11] J. Welyczko: *On Legendre curves in 3-dimensional normal almost contact metric manifolds,* Soochow J. Math., **33** (2007), no. 4, 929-937.

[12] J. Welyczko: *On Legendre curves in 3-dimensional normal almost paracontact metric manifolds,* Results Math., **54** (2009), no. 3-4, 377-387.

[13] J. C. Marrero: *The local structure of trans-Sasakian manifolds,* Ann. Mat. Pura Appl., **162** (1992), no. 4, 77-86.

[14] J. A. Oubina: *New classes of almost contact metric structures,* Publ. Math. Debrecen., **32** (1985), no 3-4, 187-193.

[15] J. I. Inoguchi: *Submanifolds with harmonic mean curvature vector eld in contact 3-manifolds,* Colloq. Math. **100** (2004), 163179.

[16] J. I. Inoguchi and J. E. Lee: *Almost contact curves in normal almost contact 3-manifolds,* J. Geom. **103** (2012), 457-474

[17] J. I. Inoguchi, J. E. Lee, *Affine biharmonic curves in 3-dimensional Homogeneous Geometries,* Mediterr. J. Math. **10** (2013), 571-592.

[18] M. Belkhelfa, I. E. Hirica and L. Verstraelen: *On legendre curves in Riemannian and Lorentzian Sasaki spaces,* Soochow J. Math. **28**(1)(2002), 81-91.

[19] S. Sasaki: *On differentiable manifolds with certain structures which are closely related to almost contact structure I,* Tôhoku Math. J., **12** (1960), 459-476.

[20] Z. Olszak: *Normal almost contact manifolds of dimension three,* Ann. Pol. Math., **47** (1986), 42-50.